

Status: Currently Official on 16-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-B0ADD591-84AB-4413-9DCF-EF10EC16B711\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3243\\_01\\_01](https://doi.org/10.31003/USPNF_R3243_01_01)  
DOI Ref: 2cc4z

© 2025 USPC  
Do not distribute

## p-Nitroaniline TS

—To 350 mg of p-nitroaniline add 1.5 mL of hydrochloric acid, and mix. Dilute with water to 50 mL, mix, and allow to settle. Place 5 mL of the clear supernatant in a 100-mL volumetric flask, and immerse it in an ice bath. While it is in the ice bath, add 1 mL of hydrochloric acid, then add, in small portions, 2 mL of sodium nitrite solution (1 in 100), dilute with water to volume, and mix.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
P-NITROANILINE TS	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocId:** [GUID-B0ADD591-84AB-4413-9DCF-EF10EC16B711\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3243\\_01\\_01](https://doi.org/10.31003/USPNF_R3243_01_01)

**DOI ref:** [2cc4z](#)

OFFICIAL