

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-A15C93BC-AE67-4866-8F7B-AB4F260B4583_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3251_01_01
DOI Ref: t1554

© 2025 USPC
Do not distribute

Phenoldisulfonic Acid TS

—Dissolve 2.5 g of phenol in 15 mL of sulfuric acid in a flask of suitable capacity. Add 7.5 mL of fuming sulfuric acid, stir well, and heat at 100° for 2 hours. Transfer the product, while still fluid, to a glass-stoppered bottle, and, when desired for use, warm in a water bath until liquefied.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
PHENOLDISULFONIC ACID TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-A15C93BC-AE67-4866-8F7B-AB4F260B4583_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3251_01_01

DOI ref: [t1554](#)

OFFICIAL