

Status: Currently Official on 16-Feb-2025

Official Date: Official Prior to 2013

Document Type: USP Monographs

DocId: GUID-52058BB1-AD58-45F5-AA06-CA01F69F2DFE_1_en-US

DOI: https://doi.org/10.31003/USPNF_M64180_01_01

DOI Ref: 7i4yh

© 2025 USPC

Do not distribute

Phenylephrine Hydrochloride Nasal Jelly

» Phenylephrine Hydrochloride Nasal Jelly contains not less than 90.0 percent and not more than 110.0 percent of the labeled amount of $C_9H_{13}NO_2 \cdot HCl$.

Packaging and storage—Preserve in tight containers.

USP REFERENCE STANDARDS (11)—

[USP Phenylephrine Hydrochloride RS](#)

Identification—Dissolve a suitable quantity in water to obtain a solution having a concentration of about 60 μ g per mL, and centrifuge, if necessary: the UV absorption spectrum of the solution so obtained exhibits maxima and minima at the same wavelengths as that of a similar solution of [USP Phenylephrine Hydrochloride RS](#), concomitantly measured.

MINIMUM FILL (755): meets the requirements.

Assay—

Mobile phase—Prepare a mixture of methanol and water (1:1) containing 1.1 g of sodium 1-octanesulfonate per liter, adjust with phosphoric acid to a pH of 3.0, filter, and degas. Make adjustments to the methanol and water ratio, if necessary (see [System Suitability](#) under [Chromatography \(621\)](#)).

Dilution solvent—Prepare a mixture of methanol and water (1:1), and adjust with phosphoric acid to a pH of 3.0.

Standard preparation—Dissolve an accurately weighed quantity of [USP Phenylephrine Hydrochloride RS](#) in **Dilution solvent** to obtain a Stock standard solution having a known concentration of about 2 mg per mL. Dilute an accurately measured volume of this solution with **Dilution solvent** to obtain the **Standard preparation** having a known concentration of about 0.1 mg per mL.

Assay preparation—Transfer an accurately weighed amount of Nasal Jelly, equivalent to about 10 mg of phenylephrine hydrochloride, to a 100-mL volumetric flask. Dilute with **Dilution solvent** to volume, and mix.

Resolution solution—Transfer 5.0 mL of Stock standard solution to a 100-mL volumetric flask, add 10 mg of [USP Epinephrine Bitartrate RS](#), dilute with **Dilution solvent** to volume, and mix.

Chromatographic system (see [Chromatography \(621\)](#))—The liquid chromatograph is equipped with a 280-nm detector and a 4.6-mm \times 25-cm column that contains packing L1. The flow rate is about 1 mL per minute. Chromatograph the **Resolution solution**: the resolution, *R*, is not less than 1.5, and the tailing factor for the phenylephrine peak is not more than 2.0. Chromatograph replicate injections of the **Standard preparation**: the relative standard deviation is not more than 2.0%.

Procedure—Separately inject equal volumes (about 20 μ L) of the **Standard preparation** and the **Assay preparation** into the chromatograph, record the chromatograms, and measure the responses for the major peaks. Calculate the quantity, in mg, of $C_9H_{13}NO_2 \cdot HCl$ in the portion of Nasal Jelly taken by the formula:

$$100C(r_u/r_s)$$

in which *C* is the concentration, in mg per mL, of [USP Phenylephrine Hydrochloride RS](#) in the **Standard preparation**, and r_u and r_s are the peak responses obtained from the **Assay preparation** and the **Standard preparation**, respectively.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
PHENYLEPHRINE HYDROCHLORIDE NASAL JELLY	Documentary Standards Support	SM22020 Small Molecules 2
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM22020 Small Molecules 2

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: GUID-52058BB1-AD58-45F5-AA06-CA01F69F2DFE_1_en-US

DOI: https://doi.org/10.31003/USPNF_M64180_01_01

DOI ref: [7i4yh](#)

OFFICIAL