

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-048B63C6-85AD-4035-97EE-7441FBEA7118_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3358_01_01
DOI Ref: t0vzh

© 2025 USPC
Do not distribute

Potassium Iodide and Starch TS:

Dissolve 0.75 g of potassium iodide in 100 mL of water. Heat to boiling, and add, with stirring, a solution of 0.5 g of soluble starch in 35 mL of water. Boil for 2 minutes, and allow to cool.

Sensitivity: Mix 15 mL in 0.05 mL of glacial acetic acid and 0.3 mL of diluted iodine TS: a blue color is produced.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
POTASSIUM IODIDE AND STARCH TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-048B63C6-85AD-4035-97EE-7441FBEA7118_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3358_01_01

DOI ref: [t0vzh](#)

OFFICIAL