

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-1D6394CF-C40D-462E-90A2-6494E5D63124_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3277_01_01
DOI Ref: 48h7r

© 2025 USPC
Do not distribute

Pyridine–Pyrazolone TS

–To 100 mL of a saturated solution of 1-phenyl-3-methyl-2-pyrazoline-5-one add 20 mL of a 1 in 1000 solution of 3,3'-dimethyl-1,1'-diphenyl-[4,4'-bi-2-pyrazoline]-5,5'-dione in pyridine. Store in a dark bottle, and use within 3 days.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
PYRIDINE–PYRAZOLONE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-1D6394CF-C40D-462E-90A2-6494E5D63124_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3277_01_01

DOI ref: [48h7r](#)

OFFICIAL