

Status: Currently Official on 16-Feb-2025  
Official Date: Official as of 01-Aug-2023  
Document Type: USP Monographs  
DocId: GUID-8858D517-C17D-49F7-9EF1-AFD57E621E2F\_4\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_M72671\\_04\\_01](https://doi.org/10.31003/USPNF_M72671_04_01)  
DOI Ref: h06m8

© 2025 USPC  
Do not distribute

## Quinidine Gluconate Extended-Release Tablets

To view the Notice from the Expert Committee that posted in conjunction with this accelerated revision, please click  
<https://www.uspnf.com/rb-quinidine-gluconate-ert-20211029>.

### DEFINITION

Quinidine Gluconate Extended-Release Tablets contain amounts of quinidine gluconate and dihydroquinidine gluconate totaling NLT 90.0% and NMT 110.0% of the labeled amount of quinidine gluconate, calculated as quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ).

### IDENTIFICATION

#### • A.

**Sample solution:** Shake an amount, equivalent to 50 mg of quinidine gluconate from powdered Tablets, with 100 mL of dilute sulfuric acid (1 in 350), and filter.

**Acceptance criteria:** The filtrate so obtained exhibits a vivid blue fluorescence when viewed under long-wavelength UV light. On the addition of hydrochloric acid, the fluorescence disappears.

#### • B.

The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

#### • C.

The  $R_F$  value of the principal spot of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in *Organic Impurities*.

### ASSAY

#### • PROCEDURE

**Solution A:** Add 35.0 mL of [methanesulfonic acid](#) to 20.0 mL of [glacial acetic acid](#), and dilute with [water](#) to 500 mL.

**Solution B:** Dissolve 10.0 mL of [diethylamine](#) in [water](#) to prepare a 100-mL solution.

**Mobile phase:** [Acetonitrile](#), *Solution A*, *Solution B*, and [water](#) (100:20:20:860). Adjust with *Solution B* to a pH of 2.6, if found to be lower.

**System suitability solution:** Transfer 10 mg each of quinidine gluconate and dihydroquinidine hydrochloride to a 50-mL volumetric flask.

Dissolve in 5 mL of [methanol](#), and dilute with *Mobile phase* to volume.

**Standard solution:** 0.2 mg/mL of [USP Quinidine Gluconate RS](#) in *Mobile phase*

**Sample stock solution:** To an amount equivalent to 160 mg of quinidine gluconate from NLT 20 finely powdered Tablets in a 100-mL volumetric flask add 80 mL of a mixture of [methanol](#) and [water](#) (1:1), and sonicate until evenly dispersed. Cool to room temperature, dilute with a mixture of [methanol](#) and [water](#) (1:1) to volume, and filter, discarding the first 20 mL of the filtrate.

**Sample solution:** 0.19 mg/mL of quinidine gluconate in *Mobile phase* prepared as follows. Transfer 3.0 mL of the *Sample stock solution* to a 25-mL volumetric flask, and dilute with *Mobile phase* to volume.

#### Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

**Mode:** LC

**Detector:** UV 235 nm

**Column:** 3- to 5-mm  $\times$  25- to 30-cm; packing [L1](#)

**Flow rate:** 1.5 mL/min

**Injection volume:** 20  $\mu$ L

#### System suitability

**Samples:** *System suitability solution* and *Standard solution*

[NOTE—The relative retention times for quinidine and dihydroquinidine are 1 and 1.5, respectively, for the *System suitability solution*.]

#### Suitability requirements

**Resolution:** NLT 1.2 between quinidine and dihydroquinidine, *System suitability solution*

**Relative standard deviation:** NMT 2.0%, *Standard solution*

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the sum of the percentages of quinidine gluconate and dihydroquinidine gluconate in the Tablets taken:

$$\text{Result} = [(r_{B,U} + r_{D,U})/(r_{B,S} + r_{D,S})] \times (C_S/C_U) \times 100$$

$r_{B,U}$  = peak response of quinidine from the *Sample solution*

$r_{D,U}$  = peak response of dihydroquinidine from the *Sample solution*

$r_{B,S}$  = peak response of quinidine from the *Standard solution*

$r_{D,S}$  = peak response of dihydroquinidine from the *Standard solution*

$C_S$  = concentration of [USP Quinidine Gluconate RS](#) in the *Standard solution* (mg/mL)

$C_U$  = nominal concentration of quinidine gluconate in the *Sample solution* (mg/mL)

**Acceptance criteria:** 90.0%–110.0%

## PERFORMANCE TESTS

### • [Dissolution \(711\)](#)

**Test 1:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 1*.

**Medium:** Add 6.9 g of [anhydrous sodium acetate](#) and 0.525 mL of [glacial acetic acid](#) to 1 L of [water](#). Adjust with 0.1 N [hydrochloric acid](#) or 0.1 N [sodium hydroxide](#) to a pH of 5.4; 900 mL.

**Apparatus 2:** 75 rpm

**Times:** 1, 2, 4, and 8 h

**Standard solution:** A known concentration of [USP Quinidine Gluconate RS](#) in *Medium*

**Sample solution:** Pass a portion of the solution under test through a suitable filter. Dilute with *Medium* if necessary, in comparison with the *Standard solution* concentration.

### Instrumental conditions

**Mode:** UV

**Analytical wavelength:** 235 nm

### Analysis

**Samples:** *Standard solution* and *Sample solution*

Determine the percentage of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved from UV absorbances of the *Sample solution* and *Standard solution*.

**Tolerances:** See [Table 1](#).

**Table 1**

Time (h)	Amount Dissolved
1	30%–50%
2	45%–65%
4	60%–85%
8	NLT 85%

The percentages of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved at the times specified conform to

[Dissolution \(711\), Acceptance Table 2](#).

**Test 4:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 4*.

**Medium:** 0.1 N [hydrochloric acid](#); 600 mL

**Apparatus 2:** 75 rpm

**Times:** Proceed as directed for *Test 1*.

**Standard solution:** A known concentration of [USP Quinidine Gluconate RS](#) in *Medium*

**Sample solution:** Pass a portion of the solution under test through a suitable filter. Dilute with *Medium* if necessary, in comparison with the *Standard solution* concentration.

### Instrumental conditions

**Mode:** UV

Analytical wavelength: 235 nm

**Analysis****Samples:** Standard solution and Sample solutionDetermine the percentage of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved from UV absorbances of the

Sample solution and Standard solution.

**Tolerances:** See [Table 2](#).**Table 2**

Time (h)	Amount Dissolved
1	30%–45%
2	45%–60%
4	60%–80%
8	NLT 85%

The percentages of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved at the times specified conform to[Dissolution \(711\), Acceptance Table 2](#).**Test 5:** If the product complies with this test, the labeling indicates that it meets USP Dissolution Test 5.**Medium:** Proceed as directed for Test 1.**Apparatus:** Proceed as directed for Test 1, using 8-mesh sinker baskets.<sup>1</sup>**Times:** 1, 2, and 4 h**Standard solution:** [USP Quinidine Gluconate RS](#) in Medium**Sample solution:** Pass a portion of the solution under test through a suitable filter. Dilute with Medium, if necessary, in comparison with the Standard solution concentration.**Instrumental conditions****Mode:** UV**Analytical wavelength:** 235 nm**Analysis****Samples:** Standard solution and Sample solutionDetermine the amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved from the UV absorbances of the Sample solution and the

Standard solution.

**Tolerances:** See [Table 3](#).**Table 3**

Time (h)	Amount Dissolved
1	20%–50%
2	40%–70%
4	NLT 75%

The percentages of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved at the times specified conform to[Dissolution \(711\), Acceptance Table 2](#).**Test 6:** If the product complies with this test, the labeling indicates that it meets USP Dissolution Test 6.**Medium:** pH 5.4 acetate buffer prepared as follows. Add 6.9 g of [anhydrous sodium acetate](#) and 0.53 mL of [glacial acetic acid](#) to 1 L of water. Adjust with 0.1 N [hydrochloric acid](#) or 0.1 N [sodium hydroxide](#) to a pH of 5.4; 900 mL, deaerated.**Apparatus 2:** 75 rpm, using 8-mesh sinker baskets<sup>1</sup>**Times:** 1, 2, 3, and 5 h

**Standard solution:** 0.012 mg/mL of [USP Quinidine Gluconate RS](#) in *Medium*

**Sample solution:** Pass a portion of the solution under test through a suitable filter. Dilute with *Medium* to a concentration similar to that of the *Standard solution*.

#### Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

**Mode:** UV

**Analytical wavelength:** 235 nm

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the concentration ( $C_i$ ) of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) in the sample withdrawn at each time point ( $i$ ):

$$\text{Result}_i = (A_U/A_S) \times C_S \times D$$

$A_U$  = absorbance of the *Sample solution*

$A_S$  = absorbance of the *Standard solution*

$C_S$  = concentration of [USP Quinidine Gluconate RS](#) in the *Standard solution* (mg/mL)

$D$  = dilution factor for the *Sample solution*

Calculate the percentage of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved at each time point ( $i$ ):

$$\text{Result}_1 = C_1 \times V \times (1/L) \times 100$$

$$\text{Result}_2 = \{[C_2 \times (V - V_S)] + [C_1 \times V_S]\} \times (1/L) \times 100$$

$$\text{Result}_3 = \{[C_3 \times [V - (2 \times V_S)]] + [(C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

$$\text{Result}_4 = \{[C_4 \times [V - (3 \times V_S)]] + [(C_3 + C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

$C_i$  = concentration of quinidine gluconate in the portion of sample withdrawn at time point ( $i$ ) (mg/mL)

$V$  = volume of *Medium*, 900 mL

$L$  = label claim (mg/Tablet)

$V_S$  = volume of the *Sample solution* withdrawn at time point ( $i$ ) (mL)

**Tolerances:** See [Table 4](#).

**Table 4**

Time point ( $i$ )	Time (h)	Amount Dissolved (%)
1	1	17–37
2	2	37–57
3	3	60–80
4	5	NLT 80

The percentages of the labeled amount of quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ) dissolved at the times specified conform to [Dissolution \(711\)](#), [Acceptance Table 2](#).

**Change to read:**

- [UNIFORMITY OF DOSAGE UNITS \(905\)](#): ▲Meet the requirements▲ (CN 1-Aug-2023)

#### Procedure for content uniformity

**Standard solution:** 0.0525 mg/mL of [USP Quinidine Gluconate RS](#) in 0.1 N hydrochloric acid

**Sample solution:** Transfer 1 intact or powdered Tablet to a 250-mL volumetric flask, and add 125 mL of 0.1 N [hydrochloric acid](#). Heat the sample with frequent agitation just to boiling, and cool to room temperature. Dilute with 0.1 N [hydrochloric acid](#) to volume, mix, and filter, discarding the first 20 mL of filtrate. If necessary, further dilute quantitatively with 0.1 N [hydrochloric acid](#).

#### Instrumental conditions

**Mode:** UV

**Cell:** 1 cm

**Analytical wavelength:** 347 nm

**Blank:** 0.1 N [hydrochloric acid](#)

#### Analysis

**Samples:** Standard solution, Sample solution, and Blank

Concomitantly determine the absorbances of the Samples.

Calculate the percentage of the labeled amounts of active ingredients, calculated as quinidine gluconate ( $C_{20}H_{24}N_2O_2 \cdot C_6H_{12}O_7$ ), in the

Tablet taken:

$$\text{Result} = (A_u/A_s) \times (C_s/C_u) \times 100$$

$A_u$  = absorbance of the Sample solution

$A_s$  = absorbance of the Standard solution

$C_s$  = concentration of [USP Quinidine Gluconate RS](#) in the Standard solution (mg/mL)

$C_u$  = nominal concentration of quinidine gluconate in the Sample solution (mg/mL)

▲ (CN 1-Aug-2023)

#### IMPURITIES

##### • ORGANIC IMPURITIES

**Standard solution A:** 6 mg/mL of [USP Quinidine Gluconate RS](#) in diluted alcohol

**Standard solution B:** 0.06 mg/mL of [USP Quinidine Gluconate RS](#) in diluted alcohol from Standard solution A

**Standard solution C:** 0.04 mg/mL of [USP Quinonone RS](#) (corresponding to 0.06 mg of the gluconate) in diluted alcohol

**Sample solution:** Nominally equivalent to 6 mg/mL of quinidine gluconate prepared as follows. Shake a quantity of powdered Tablets, equivalent to about 150 mg of quinidine gluconate, with 25 mL of diluted alcohol for 10 min, and filter.

#### Chromatographic system

(See [Chromatography \(621\), General Procedures, Thin-Layer Chromatography.](#))

**Mode:** TLC

**Adsorbent:** 0.25-mm layer of chromatographic silica gel mixture

**Application volume:** 10  $\mu$ L

**Developing solvent system:** Chloroform, acetone, and diethylamine (50:40:10)

#### Analysis

**Samples:** Standard solution A, Standard solution B, Standard solution C, and Sample solution

Proceed as directed in the chapter. The solvent chamber is used without previous equilibration. When the solvent front has moved 15 cm, remove the plate from the chamber, mark the solvent front, and allow the solvent to evaporate. Spray with glacial acetic acid. Locate the spots on the plate by examination under long-wavelength UV light.

**Acceptance criteria:** Any spot produced by the Sample solution at the  $R_f$  value of a spot produced by Standard solution C is not greater in size or intensity than that corresponding spot. Apart from these spots and from the spots appearing at the  $R_f$  value of quinidine gluconate and dihydroquinidine gluconate (the two spots most evident from Standard solution A), any additional fluorescent spot is not greater in size or intensity than the principal spot of Standard solution B.

#### ADDITIONAL REQUIREMENTS

• **PACKAGING AND STORAGE:** Preserve in well-closed, light-resistant containers.

• **LABELING:** The labeling indicates the *Dissolution Test* with which the product complies.

• [USP REFERENCE STANDARDS \(11\)](#)

[USP Quinidine Gluconate RS](#)

[USP Quinonone RS](#)

Cinchonan-9-one, 6'-methoxy-, (8 $\alpha$ )-

$C_{20}H_{22}N_2O_2$  322.40

<sup>1</sup> A suitable sinker is available from [www.agilent.com](http://www.agilent.com), catalog number 12-3062.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
QUINIDINE GLUCONATE EXTENDED-RELEASE TABLETS	<a href="#">Documentary Standards Support</a>	SM22020 Small Molecules 2
REFERENCE STANDARD SUPPORT	RS Technical Services <a href="mailto:RSTECH@usp.org">RSTECH@usp.org</a>	SM22020 Small Molecules 2

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 31(1)

**Current DocID: GUID-8858D517-C17D-49F7-9EF1-AFD57E621E2F\_4\_en-US**

**DOI: [https://doi.org/10.31003/USPNF\\_M72671\\_04\\_01](https://doi.org/10.31003/USPNF_M72671_04_01)**

**DOI ref: h06m8**

OFFICIAL