

Rectiofar

rectal solution



PHARMEDIC

COMPOSITION

- Glycerin59,53 g
- Purified watersqf 100 ml.

INDICATIONS

Treatment of constipation in children and adults.

DOSAGE AND ADMINISTRATION:

Inject drug into the rectum, 2 tubes daily, 1 – 2 times daily.

- Children: 3 ml.
- Adults: 5 ml.

CONTRAINDICATIONS

- Hypersensitivity to any of the components of the drug.
- Acute hemorrhoids, anal fistula, diarrhea.

PRECAUTIONS

- Patients with cardiac, renal, hepatic diseases.
- Elderly.
- Patients with confused mental state, congestive heart failure, senescence, diabetes and severe dehydration.

INTERACTIONS: none.

USE IN PREGNANCY AND LACTATION

- Glycerin can be used during pregnancy when clearly needed.
- Drug should be used with caution in nursing women.

EFFECTS ON ABILITY TO DRIVE OR OPERATE MACHINERY None

ADVERSE REACTIONS

- Glycerin can cause irritation when given rectally.
 - After rectal administration, the absorption of drug is poor but when absorbed:
 - Drug may cause headache, vomiting.
 - Rare cases of dizziness, confusion, diarrhea, nausea, thirst.
 - Some symptoms may occur very rarely: arrhythmias, hyperglycemia.
- * **Contact your physician if you experience any adverse reactions while using this medicine.**

OVERDOSAGE

- Overdosage may cause excessive diarrhoea, nausea, arrhythmias, rectal irritation, burning, cramping pain and hyperglycemia.
- With overdosage, discontinue the drug and hospitalize the patient.

PHARMACODYNAMICS

Glycerin is administered rectally to draw fluid into the colon and thus promotes fecal evacuation. It may also have lubricating and fecal softening actions.

PHARMACOKINETICS

When rectal administration, the absorption of drug is poor.

EXTERNAL USE ONLY - DO NOT SWALLOW.

KEEP OUT OF REACH OF CHILDREN

READ THE LEAFLET CAREFULLY BEFORE USE.

FOR MORE INFORMATION, CONSULT YOUR PHYSICIAN.

SHELF - LIFE: 30 months from date of manufacturing.

SPECIFICATION: Manufacturer's

PRESENTATION

- Box of 50 sachets of 1 tube (attached plate) of 3 ml.
- Box of 50 sachets of 1 tube of 3 ml
- Box of 40 sachets of 1 tube (attached plate) of 5 ml.
- Box of 40 sachets of 1 tube of 5 ml.

DO NOT STORE OVER 30°C.

PHARMEDIC JSC: 367 Nguyen Trai Street, District 1, Ho Chi Minh City, Vietnam.

Manufactured by PHARMEDIC in compliance with GMP - WHO norms
1/67 Nguyen Van Qua Street, District 12, Ho Chi Minh City, Vietnam.