

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-9E85D506-47CC-4095-8B10-18CF866D12F8_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3300_01_01
DOI Ref: rrc6f

© 2025 USPC
Do not distribute

Sodium Fluoride TS

—Dry about 500 mg of sodium fluoride at 200° for 4 hours. Accurately weigh 222 mg of the dried material, and dissolve in water to make 100.0 mL. Pipet 10 mL of this solution into a 1-L volumetric flask, and dilute with water to volume. Each mL of this solution corresponds to 0.01 mg of fluorine (F).

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
SODIUM FLUORIDE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-9E85D506-47CC-4095-8B10-18CF866D12F8_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3300_01_01

DOI ref: [rrc6f](#)

OFFICIAL