

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-DDD28495-6E2E-4A34-BBDB-60AE8937CF5E_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3305_01_01
DOI Ref: k12cm

© 2025 USPC
Do not distribute

Sodium Iodohydroxyquinolinesulfonate TS

—Dissolve 8.8 g of iodohydroxyquinoline sulfonic acid in 200 mL of water, and add 6.5 mL of 4 N sodium hydroxide. Dilute with water to 250 mL, mix, and filter.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
SODIUM IODOHYDROXYQUINOLINESULFONATE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-DDD28495-6E2E-4A34-BBDB-60AE8937CF5E_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3305_01_01

DOI ref: [k12cm](#)

OFFICIAL