

Status: Currently Official on 16-Feb-2025
 Official Date: Official as of 01-May-2021
 Document Type: USP Monographs
 DocId: GUID-CEE865C8-04DE-43ED-A71F-1965FA341932_2_en-US
 DOI: https://doi.org/10.31003/USPNF_M77004_02_01
 DOI Ref: 7zh9q

© 2025 USPC
 Do not distribute

Add the following:

^Sodium Phenylbutyrate Tablets

DEFINITION

Sodium Phenylbutyrate Tablets contain NLT 90.0% and NMT 110.0% of the labeled amount of sodium phenylbutyrate ($C_{10}H_{11}NaO_2$).

IDENTIFICATION

- **A.** The UV spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.
- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• PROCEDURE

Mobile phase: [Methanol](#), [water](#), and [glacial acetic acid](#) (49:50:1)

Diluent: [Methanol](#) and [water](#) (20:80)

Standard solution: 2.5 mg/mL of [USP Sodium Phenylbutyrate RS](#) in *Diluent*

Sample solution: Nominally 2.5 mg/mL of sodium phenylbutyrate prepared as follows. Finely powder NLT 10 Tablets and transfer a suitable portion equivalent to about 500 mg of sodium phenylbutyrate to a 200-mL volumetric flask, add about 140 mL of *Diluent*, and sonicate for 10 min with intermittent shaking. Dilute with *Diluent* to volume and mix well. Pass through a suitable filter of 0.45- μ m pore size.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 260 nm. For *Identification A*, use a diode array detector in the range of 200–400 nm.

Column: 4.6-mm \times 15-cm; 5- μ m packing [L1](#)

Flow rate: 1.2 mL/min

Injection volume: 20 μ L

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of sodium phenylbutyrate ($C_{10}H_{11}NaO_2$) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Sodium Phenylbutyrate RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of sodium phenylbutyrate in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

• [DISSOLUTION \(711\)](#)

Medium: [Simulated intestinal fluid](#) without enzyme; 900 mL

Apparatus 2: 75 rpm**Time:** 30 min**Standard solution:** ($L/900$) mg/mL of [USP Sodium Phenylbutyrate RS](#) in *Medium*, where L is the label claim in mg/Tablet. Sonicate, if necessary, to dissolve.**Sample solution:** Pass a portion of the solution under test through a suitable filter of 0.45- μm pore size.**Instrumental conditions****Mode:** UV**Analytical wavelength:** 260 nm**Analysis****Samples:** *Standard solution* and *Sample solution*Calculate the percentage of the labeled amount of sodium phenylbutyrate ($\text{C}_{10}\text{H}_{11}\text{NaO}_2$) dissolved:

$$\text{Result} = (A_u/A_s) \times (C_s/L) \times V \times 100$$

 A_u = absorbance of the *Sample solution* A_s = absorbance of the *Standard solution* C_s = concentration of [USP Sodium Phenylbutyrate RS](#) in the *Standard solution* (mg/mL) L = label claim (mg/Tablet) V = volume of *Medium*, 900 mL**Tolerances:** NLT 85% (Q) of the labeled amount of sodium phenylbutyrate ($\text{C}_{10}\text{H}_{11}\text{NaO}_2$) is dissolved.

- [Uniformity of Dosage Units \(905\)](#): Meet the requirements

IMPURITIES**• ORGANIC IMPURITIES****Mobile phase:** [Methanol](#), [water](#), and [glacial acetic acid](#) (44:55:1)**Diluent:** [Methanol](#) and [water](#) (20:80)**System suitability stock solution A:** 0.08 mg/mL of [USP Phenylbutyrate Related Compound A RS](#) in *Diluent*. Sonicate to dissolve, if necessary.**System suitability stock solution B:** 0.06 mg/mL of [USP Phenylbutyrate Related Compound B RS](#) in *Diluent*. Sonicate to dissolve, if necessary.**System suitability solution:** 4 mg/mL of [USP Sodium Phenylbutyrate RS](#), 0.008 mg/mL of [USP Phenylbutyrate Related Compound A RS](#), and 0.006 mg/mL of [USP Phenylbutyrate Related Compound B RS](#) prepared as follows. Dissolve a suitable amount of [USP Sodium Phenylbutyrate RS](#) with 50% of the total flask volume of [methanol](#) and sonicate, if necessary, to dissolve. Add suitable volumes of *System suitability stock solution A* and *System suitability stock solution B* to the same flask and dilute with *Diluent* to volume.**Sensitivity solution:** 1.5 $\mu\text{g}/\text{mL}$ each of [USP Phenylbutyrate Related Compound A RS](#) and [USP Phenylbutyrate Related Compound B RS](#) from *System suitability stock solution A* and *System suitability stock solution B* in *Diluent***Standard solution:** 0.008 mg/mL of [USP Phenylbutyrate Related Compound A RS](#) and 0.006 mg/mL of [USP Phenylbutyrate Related Compound B RS](#) from *System suitability stock solution A* and *System suitability stock solution B* in *Diluent***Sample solution:** Nominally 4 mg/mL of sodium phenylbutyrate in *Diluent* prepared as follows. Transfer a suitable quantity from NLT 20 powdered Tablets, equivalent to about 200 mg of sodium phenylbutyrate, to a 50-mL volumetric flask. Add about 35 mL of *Diluent* and sonicate with intermittent shaking. Dilute with *Diluent* to volume. Pass through a suitable filter of 0.45- μm pore size.**Chromatographic system**(See [Chromatography \(621\), System Suitability](#).)**Mode:** LC**Detector:** UV 245 nm**Column:** 2.1-mm \times 10-cm; 1.7- μm packing [L1](#)**Column temperature:** 35°**Flow rate:** 0.2 mL/min**Injection volume:** 2 μL **System suitability****Samples:** *System suitability solution*, *Sensitivity solution*, and *Standard solution***Suitability requirements****Resolution:** NLT 6.0 between the phenylbutyrate related compound B and phenylbutyrate peaks, *System suitability solution*

Relative standard deviation: NMT 5.0% for the phenylbutyrate related compound A and phenylbutyrate related compound B peaks,

Standard solution

Signal-to-noise ratio: NLT 10 for the phenylbutyrate related compound A and phenylbutyrate related compound B peaks, *Sensitivity solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of phenylbutyrate related compound A or phenylbutyrate related compound B in the portion of Tablets taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

r_u = peak response of phenylbutyrate related compound A or phenylbutyrate related compound B from the *Sample solution*

r_s = peak response of phenylbutyrate related compound A or phenylbutyrate related compound B from the *Standard solution*

C_s = concentration of [USP Phenylbutyrate Related Compound A RS](#) or [USP Phenylbutyrate Related Compound B RS](#) in the *Standard solution* (mg/mL)

C_u = nominal concentration of sodium phenylbutyrate in the *Sample solution* (mg/mL)

Calculate the percentage of any individual unspecified impurity in the portion of Tablets taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

r_u = peak response of any individual unspecified impurity from the *Sample solution*

r_s = peak response of phenylbutyrate related compound A from the *Standard solution*

C_s = concentration of [USP Phenylbutyrate Related Compound A RS](#) in the *Standard solution* (mg/mL)

C_u = nominal concentration of sodium phenylbutyrate in the *Sample solution* (mg/mL)

Acceptance criteria: See [Table 1](#). The reporting threshold is 0.05%, except for phenylbutyrate related compound B.

Table 1

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Phenylbutyrate related compound A	0.30	0.10
Phenylbutyrate related compound B	0.68	0.05
Sodium phenylbutyrate	1.0	—
Any individual unspecified impurity	—	0.10
Total impurities	—	0.5

ADDITIONAL REQUIREMENTS

• **PACKAGING AND STORAGE:** Preserve in tight containers. Store at controlled room temperature.

• **USP REFERENCE STANDARDS (11).**

[USP Sodium Phenylbutyrate RS](#)

[USP Phenylbutyrate Related Compound A RS](#)

3-Benzoylpropionic acid;

4-Oxo-4-phenylbutanoic acid.

$C_{10}H_{10}O_3$ 178.18

[USP Phenylbutyrate Related Compound B RS](#)

α -Tetralone;

3,4-Dihydronaphthalen-1(2H)-one.

$C_{10}H_{10}O$ 146.19 ▲ (USP 1-May-2021)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
SODIUM PHENYLBUTYRATE TABLETS	Documentary Standards Support	SM32020 Small Molecules 3
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 44(2)

Current DocID: [GUID-CEE865C8-04DE-43ED-A71F-1965FA341932_2_en-US](#)

DOI: https://doi.org/10.31003/USPNF_M77004_02_01

DOI ref: [7zh9q](#)

<https://trungtamthuoc.com/>