

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-7352B3C7-EF67-443B-B941-36235AC63BEA_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3327_01_01
DOI Ref: tyd4I

© 2025 USPC
Do not distribute

Sulfanilic-1-Naphthylamine TS

—Dissolve 500 mg of sulfanilic acid in 150 mL of acetic acid. Dissolve 100 mg of 1-naphthylamine hydrochloride in 150 mL of acetic acid, and mix the two solutions. The pink color that may develop on standing can be removed by treatment with zinc.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
SULFANILIC-1-NAPHTHYLAMINE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-7352B3C7-EF67-443B-B941-36235AC63BEA_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3327_01_01

DOI ref: [tyd4I](#)

OFFICIAL