

Status: Currently Official on 16-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-4B13912E-C7E3-4FB2-94F6-3A91DD5B4743_1_en-US
DOI: https://doi.org/10.31003/USPNF_R1541_01_01
DOI Ref: l7num

© 2025 USPC
Do not distribute

Tetraethylene Glycol,

$C_8H_{18}O_5$ 194.23 CAS RN[®]: 112-60-7.—Nearly colorless liquid. Refractive index: about 1.46.

Assay: When examined by gas–liquid chromatography, using suitable gas chromatographic apparatus and conditions, it shows a purity of not less than 90%.

Boiling Range (Reagent test): between 177° and 187°, at a pressure of 9 mm of Hg.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
TETRAETHYLENE GLYCOL	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-4B13912E-C7E3-4FB2-94F6-3A91DD5B4743_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R1541_01_01

DOI ref: [l7num](#)