

Status: Currently Official on 16-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: USP Monographs  
DocId: GUID-EDB63F90-A1EF-488C-9A15-0734A7226B5C\_1\_en-US  
DOI: https://doi.org/10.31003/USPNF\_M82480\_01\_01  
DOI Ref: 06y3c

© 2025 USPC  
Do not distribute

# Thiamine Hydrochloride Oral Solution

## DEFINITION

Thiamine Hydrochloride Oral Solution contains NLT 95.0% and NMT 135.0% of the labeled quantity of thiamine hydrochloride ( $C_{12}H_{17}ClN_4OS \cdot HCl$ ).

## IDENTIFICATION

### A.

**Sample solution:** Dilute a portion of Oral Solution with water to a concentration of 10 mg/mL of thiamine hydrochloride.

**Analysis:** To 0.5 mL of the *Sample solution* add 5 mL of 0.5 N sodium hydroxide, then add 0.5 mL of potassium ferricyanide TS and 5 mL of isobutyl alcohol. Shake the mixture vigorously for 2 min, and allow the liquid layers to separate. Illuminate from above by a vertical beam of UV light, and observe the air–liquid meniscus at a right angle to this beam.

**Acceptance criteria:** The air–liquid meniscus shows a vivid blue fluorescence, which disappears when the mixture is slightly acidified, but reappears when it is again made alkaline.

## ASSAY

### PROCEDURE

**Mobile phase:** Methanol and 0.04 M aqueous monobasic potassium phosphate (45:55)

**Internal standard solution:** 100 µg/mL of methylparaben in *Mobile phase*

**Standard stock solution:** 500 µg/mL of [USP Thiamine Hydrochloride RS](#) in *Mobile phase*

**Standard solution:** Dilute a mixture of equal volumes of the *Standard stock solution* and *Internal standard solution* with *Mobile phase* to obtain a concentration of [USP Thiamine Hydrochloride RS](#) of about 50 µg/mL.

**Sample stock solution:** Equivalent to 500 µg/mL of thiamine hydrochloride in *Mobile phase* from an accurately measured volume of Oral Solution

**Sample solution:** Dilute a mixture of equal volumes of the *Sample stock solution* and *Internal standard solution* with *Mobile phase* to obtain a concentration of thiamine hydrochloride of about 50 µg/mL.

### Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** LC

**Detector:** UV 254 nm

**Column:** 3.9-mm × 30-cm; packing L1

**Flow rate:** 1 mL/min

**Injection size:** 25 µL

### System suitability

**Sample:** *Standard solution*

[NOTE—The relative retention times for thiamine and methylparaben are about 0.35 and 1.0, respectively.]

### Suitability requirements

**Resolution:** NLT 6.0 between thiamine and methylparaben

**Relative standard deviation:** NMT 2.0%

### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of thiamine hydrochloride ( $C_{12}H_{17}ClN_4OS \cdot HCl$ ) in the portion of Oral Solution taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

$R_U$  = peak area ratio of thiamine to methylparaben from the *Sample solution*

$R_s$  = peak area ratio of thiamine to methylparaben from the *Standard solution*

$C_s$  = concentration of [USP Thiamine Hydrochloride RS](#) in the *Standard stock solution* (µg/mL)

$C_u$  = nominal concentration of thiamine hydrochloride in the *Sample stock solution* (µg/mL)

**Acceptance criteria:** 95.0%–135.0%

**OTHER COMPONENTS**

- **ALCOHOL DETERMINATION, Method II(611):** 90.0%–110.0% of the labeled quantity of C<sub>2</sub>H<sub>5</sub>OH, using acetone as the internal standard

**ADDITIONAL REQUIREMENTS**

- **PACKAGING AND STORAGE:** Preserve in tight, light-resistant containers.
- **USP REFERENCE STANDARDS (11):**  
[USP Thiamine Hydrochloride RS](#)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
THIAMINE HYDROCHLORIDE ORAL SOLUTION	<a href="#">Natalia Davydova</a> Scientific Liaison	NBDS2020 Non-botanical Dietary Supplements
REFERENCE STANDARD SUPPORT	RS Technical Services <a href="mailto:RSTECH@usp.org">RSTECH@usp.org</a>	NBDS2020 Non-botanical Dietary Supplements

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 28(2)

**Current DocID:** GUID-EDB63F90-A1EF-488C-9A15-0734A7226B5C\_1\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_M82480\\_01\\_01](https://doi.org/10.31003/USPNF_M82480_01_01)

**DOI ref:** [06y3c](#)

