

CEELIN Sirô uống dạng giọt
Hộp và nhǎn chai 15 mL

3/4/14/0


BỘ Y TẾ
CỤC QUẢN LÝ DƯỢC
ĐÃ PHÊ DUYỆT

Lần đầu: 19/6/2013

Label 15 mL: 64 x 32 mm



Box 15 mL: 52 x 28 x 100 mm



<https://trungtamthuoc.com/>



© Trung tâm Dược

Sản xuất tại CÔNG TY TNHH UNITED INTERNATIONAL PHARMA
39/10A LAMP, 19/10A Phan Văn Trị, Phường 11, Quận Gò Vấp, TP.HCM
Thành Phố Hồ Chí Minh, Việt Nam - Singapore, Phòng Hoa Phu
Thị trấn Mai Mỹ, Bình Dương Province, Vietnam Tel: 0819621000

© Registered trademarks of
Pediatrics, Inc.



CEELIN® Vitamin C

100 mg/ mL Drops - 100 mg/ 5 mL Syrup

SUPPORT RESISTANCE TO INFECTION AND IMMUNE FUNCTION

PHARMACOLOGY – BENEFIT OF VITAMIN C

CEELIN® (vitamin C) involved in metabolism of phenylalanine, tyrosine, folic acid, norepinephrine, histamine, iron, and some drug enzyme systems, utilization of carbohydrates, synthesis of lipids, proteins, immune function, resistance to bacterial infections and preservation of blood vessel integrity. Vitamin C deficiency caused of scurvy disease.

CEELIN® (vitamin C) is necessary for collagen formation, tissue repair in the body and involved in some oxidation-reduction reactions, helps formation and maintenance of firm and healthy gums, teeth, cartilages, skeletal, vascular, and connective tissues, wound healing.

COMPOSITION

Each 1 mL CEELIN® Drops contains:

Vitamin C (as Sodium Ascorbate) 100 mg

Excipients: EDTA, Sodium Metabisulfite, Methyl Paraben, Propyl Paraben, Glycerin, Sorbitol Solution,

Sucralose, Sodium Chloride, Propylene Glycol, D&C Yellow, D&C Red, Orange Juice Flavor, Citric Acid,

Purified Water.

Each 5 mL (1 teaspoon) CEELIN® Syrup contains:

Vitamin C (as Sodium Ascorbate) 100 mg

Excipients: EDTA, Sodium Metabisulfite, Methyl Paraben, Propyl Paraben, Mannitol, Sorbitol Solution,

Sucralose, Sodium Chloride, Propylene Glycol, D&C Yellow, D&C Red, Orange Juice Flavor, Citric Acid,

Purified Water.

INDICATIONS

For the prevention and treatment of vitamin C deficiency in infants and children. Support resistance to infection and immune function.

DOSE AND INSTRUCTION FOR USE

Orally. Prevention and therapeutic dose per day:

	CEELIN® 100 mg/ mL Drops	CEELIN® 100 mg/ 5 mL Syrup
Prevention	0.25 – 0.5 mL (5 - 10 drops)/ time 1 time/ day	2.5 mL (1/2 teaspoon)/ time 1 time/ day
Therapeutic	1 mL (20 drops)/ time 1-3 times/ day	5 mL (1 teaspoon)/ time 1-3 times/ day

Or as prescribed by the physician.

CONTRAINDICATIONS

Contraindication taking high doses of vitamin C in the person who had G6PD deficiency. Hypersensitivity to vitamin C, Thalassemia, history of renal stones, hyperoxaluria and oxalate metabolism disorders.

PRECAUTION

Vitamin C should be given with care to patients with hyperoxaluria. Tolerance may be induced in patients taking high doses.

Effects on ability to drive and use machines: the status of drowsiness had occurred.

PREGNANCY AND LACTATION

Pregnancy: vitamin C crosses the placenta. Studies have not been done in humans and animals. Problems in humans have not been documented with intake of normal daily recommended amounts. However, ingestion of large quantities of vitamin C daily throughout pregnancy may increase the require of vitamin C and cause of scurvy disease in newborn. Lactation: vitamin C is distributed into breast milk. Problems in humans and the infants have not been documented with intake of normal daily recommended amounts.

ADVERSE DRUG REACTIONS (ADR)

Hyperoxaluria, nausea or vomiting, heartburn, abdominal muscle spasms, fatigue, flushing or redness of skin, headache, insomnia and the status of drowsiness had occurred. Oral doses greater than 1 gram per day can cause of diarrhea.

Frequently, ADR > 1/100

Renal: hyperoxaluria.

Rarely, 1/1000 < ADR < 1/100

Blood: haemolysis anemia.

Cardiovascular: flushing, heart failure.

Gastro – intestinal: nausea, vomiting, abdominal pain, abdominal muscle spasms, flatulence, diarrhea.

Nerve – muscles and bones: flank pain.

Please inform your doctor of all undesirable effects upon drug administration.

OVERDOSE AND TREATMENT

The symptoms of vitamin C overdose include kidney stones, nausea, gastritis and diarrhea. The diuretic therapy by intravenous infusion may be effective in cases of excessive dose of oral vitamin C.

DRUG INTERACTIONS

Vitamin C can significantly enhance iron uptake and metabolism.

Concurrent use of acetylsalicylic acid (aspirin) with vitamin C may increase excretion of vitamin C & reduce excretion of acetylsalicylic acid in urine.

PRESENTATION

CEELIN® 100 mg/ mL Drops – Box of 15 mL bottle.

CEELIN® 100 mg/ 5 mL Syrup – Box of 30 mL,
60 mL, 120 mL bottle.

STORAGE

Always keep container tightly closed.

Store at temperatures not exceeding 30°C.

SHELF-LIFE

24 months from manufacturing date.

KEEP OUT OF REACH OF CHILDREN


READ THE PACKAGE INSERT CAREFULLY BEFORE USE

FOR MORE INFORMATION, PLEASE ASK FOR DOCTOR'S ADVICE

Manufactured by UNITED INTERNATIONAL PHARMA CO., LTD.

WHO-GMP, GLP, GSP, No.16 VSIP II, Street No.7, Vietnam-Singapore II Industrial Park,
Hoa Phu Ward, Thu Dau Mot City, Binh Duong Province, Vietnam. Tel: 08-39621000

® Registered trademark of Pediatrica, Inc.



INVIE02/7/00010-0-04