

Status: Currently Official on 17-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-BF1D2021-6C8E-4158-8612-ADC704BEEF8E\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3342\\_01\\_01](https://doi.org/10.31003/USPNF_R3342_01_01)  
DOI Ref: s1bid

© 2025 USPC  
Do not distribute

## Titanium Trichloride–Sulfuric Acid TS

–Mix carefully 20 mL of titanium trichloride TS in 13 mL of sulfuric acid. Add sufficient 30% hydrogen peroxide to produce a yellow color. Heat until white fumes are evolved, allow to cool, and dilute with water. Repeat the evaporation and addition of water until a colorless solution is obtained. Dilute with water to 100 mL.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
TITANIUM TRICHLORIDE–SULFURIC ACID TS	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocId:** [GUID-BF1D2021-6C8E-4158-8612-ADC704BEEF8E\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3342\\_01\\_01](https://doi.org/10.31003/USPNF_R3342_01_01)

**DOI ref:** [s1bid](#)

OFFICIAL