Clindamycin & Clotrimazole Soft Gelatin Vaginal Capsules

Tivag

Mỗi viên nang âm đạo Gelatin mềm chứa:

Clindamycin Phosphat	BP	
Phương trình đến Clindamycin		100 mg
Clotrimazole	BP	200 mg
Tá dược vừa đủ		

Màu được phê duyệt được sử dụng trong vỏ viên nang Gelatin mềm.

Cơ chế tác dụng của Tivag

Các thành phần chính có trong 1 viên đặt âm đạo **Tivag.**

Clotrimazole: Clotrimazole có tính kháng nấm phổ rộng, tác động tốt lên tính thấm thấu của vách tế bào vi nấm nhờ khả năng can thiệt lên các lipid của màng.Clotrimazole được bài tiết ở đường tiểu và đường mật chỉ với khoảng dưới dạng hoạt động.

Clindamycin: Đây là một loại kháng sinh thuộc họ Lincosamid có tác dụng ức chế tổng hợp Protein của vi khuẩn nhờ vào khả năng liên kết với tiểu phần 50S của Ribosom. Khoảng hơn Clindamycin được dung nạp vào cơ thể sẽ liên kết với Protein huyết tương. Hoạt chất này thường được chuyển hóa chủ yếu ở gan, thời gian bán thải của Clindamycin ở vào khoảng 2 - 3 giờ đồng hồ.

Chỉ định :

- **Tivag** thường được chỉ định trong điều trị viêm âm đạo, âm hỗ hỗn hợp do một số loại vi khuẩn, nấm gây nên như: Gardnerellavaginalis, Mycoplasma sp, Candida albicans, Mobilicus.
- Điều trị các nhiễm khuẩn do vi khuẩn kỵ khí.
- Nhiễm khuẩn phụ khoa: Viêm nội mạc tử cung , viêm cơ nội mạc tử cung, áp xe vòi buồng trứng.
- Viêm âm đạo không đặc hiệu.
- Nhiễm Trichomonassinhdục.
- Ngoài ra, **Tivag** còn có thể được bác sĩ chỉ định dùng trong các trường hợp khác.Đừng tự ý mua **Tivag** về dùng cho bất cứ mục đích nào khi bạn chưa nhận được tham vấn y khoa.

Chống chỉ định : Tivag chống chỉ định trong các trường hợp đề cập dưới đây:

- Đối tượng quá mẫn với thành phần có trong **Tivag.**
- Loạn tạo máu hoặc có tiền sử loạn chuyển hóa porphyrin cấp.
- Người bệnh có các rối loạn thần kinh thực thể.

THẬN TRỌNG:

Nên cẩn trọng khi dùng **Tivag** với phụ nữ mang thai, nhất là trong 3 tháng đầu của thai kỳ.Nếu thật sự cần thiết hãy tham khảo ý kiến của Bác sỹ trước khi sử dụng. Chưa có bằng chứng **Tivag** ảnh hưởng có hại đến giai đoạn của thai kỳ.Cần phải cân nhắc giữa lợi ích khi dùng **Tivag** với khả năng gây hại cho thai nhi.Chưa có thông tin chính thức về việc **Tivag** có tác động đến sữa mẹ hay không, tuy nhiên phụ nữ đang cho con bú chỉ sử dụng khi có chỉ định của Bác sỹ.

Tivag dùng để đặt âm đạo, tuyệt đối không sử dụng bằng đường uống. Trường hợp không may nuốt phải, cần báo ngay cho Bác sỹ để xử lý kịp thời. Không nên quan hệ tình dục trong suốt quá trình điều trị bằng **Tivag**. Để đề phòng tái nhiễm, cần chú ý điều trị cho cả bạn tình.

Cách Dùng :

- Trước khi đặt **Tivag** , bạn cần vệ sinh vùng kín sạch sẽ. Có thể dùng nước muối loãng để vệ sinh, tránh dùng xà bông hay chất tẩy rửa. Trường hợp muốn dùng dung dịch vệ sinh chuyên dụng, bạn cần có được sự cho phép của bác sĩ.
- Cách đặt: Nằm ngửa ở tư thế đầu gối co lên, hai chân dạng ra. Dùng 2 ngón tay kẹp chặt viên **Tivag** rồi đặt vào âm đạo. Sau đó dùng 1 ngon tay để đẩy viên **Tivag** càng sâu càng tốt. Kết thúc việc đặt **Tivag**, bạn khép chân lại và nằm nghỉ ít nhất
  Cần vệ sinh tay sạch sẽ cả trước và sau khi tiếp xúc với **Tivag**. Cắt gọn móng hay, có thể đeo bao cao su cho ngón tay khi đặt **Tivag** để đảm bảo an toàn và tránh làm tổn thương âm đạo.

Liều lượng : Theo chỉ định của Bác sỹ: Liều lượng và tần suất dùng **Tivag** được đề cập dưới đây chỉ mang tính tham khảo, đáp ứng với các trường hợp thường gặp.

Liều phổ biến: 1 viên/lần/ngày, vào buổi tối trước khi đi ngủ.

Trường hợp bệnh nặng: 1 viên/lần, 2 lần/ngày, vào buổi sáng và buổi tối.

Tuyệt đối không tự ý tăng, giảm liều hoặc ngưng dùng trong bất cứ trường hợp nào. Khi liều dùng không đáp ứng, nên báo ngay cho bác sĩ để có biện pháp khắc phục kịp thời.

Lưu ý: Tác dụng của **Tivag** tương đối mạnh, vì thế sau sử dụng nếu không thấy triệu chứng thuyên giảm hãy báo ngay cho bác sĩ.

Có thể bạn sẽ được chỉ định thay thế bằng loại thuốc khác phù hợp hơn.

Bảo quản Tivag : Bạn có thể bảo quản **Tivag** theo hướng dẫn dưới đây để đảm bảo tác dụng điều trị:

Giữ **Tivag** ở nhiệt độ phòng ở khoản tốt nhất là

Để nơi khô thoáng, tránh ánh sáng trực tiếp và độ ẩm vượt

Tránh xa tầm tay với của trẻ em.

Chỉ sử dụng để đặt âm đạo, không dùng uống.

ĐIỀU KIỆN BẢO QUẢN: Bảo quản dưới 25 ° C ở nơi khô mát. Bảo vệ khỏi ánh nắng trực tiếp, nhiệt và độ ẩm.

Mfg Lic. No: 82/UA/2017
Manufactured. by :
Renowed Life Sciences
WHO-GMP Certified Co.
Plot No. 10 & 11, Sector 6-B, IIE, SIDCUL,
Haridwar 249403 , Uttarakhand (INDIA)

https://trungtamthuoc.com/

Clindamycin & Clotrimazole Soft Gelatin Vaginal Capsules

Tivag

Each Soft Gelatin Vaginal Capsule Contains:

Clindamycin Phosphate	BP	
Eq. to Clindamycin		100 mg
Clotrimazole	BP	200 mg
Excipients		q.s.

Approved Colour used in Soft Gelatin capsule shells.

Properties: Clotrimazole is an imidazole and has broad spectrum anti fungal action. It brings about its therapeutic effect by inhibiting ergo sterol synthesis. Inhibition of ergo sterol synthesis alters the permeability of the cell membrane of sensitive fungi and leads to structural and functional impairment of the cytoplasmic membrane. It has fungicidal action against *Candida albicans*, other species of Candida as well as *tricomonacidal* action.
Clindamycin is a lincosamide antibiotic primarily acting against Gram positive and wide range of anaerobic bacteria. It is congener of lincomycin. Clindamycin inhibits bacterial protein synthesis by its action at the bacterial ribosome. The antibiotic binds preferentially to the 50s ribosomal subunits and affects the process of protein synthesis. Clindamycin has antimicrobial activity against bacteroides spp., *Gerdnerella vaginalis, Mobiluncus spp., Mycoplasma hominis, Peptostreptococcus spp., Haemophilus spp.*, and *Corynrbacterium.*

Indications: Infective Leucorrhoea, mixed infection and non specific vaginitis. Vaginal candidiasis, Bacterial vaginosis, Trichomoniasis, Cadida vulvitis & vaginal mycoses.

Contraindication: Tivag is a contraindicated in women with a history of hypersensitivity to clotrimazole (or azole group) and Clindamycin (or lincomycin group). Tivag also contraindicated in women with a history of regional enteritis, ulcerative colitis or "antibiotic associated" colitis (pseudo membranous colitis).

Warnings and precautions: The patient should be instructed not to engage in vaginal intercourse or use of other vaginal products (such as tampons or douches) during treatment with clindamycin.
If there is a lack of response to Tivag, appropriate microbiological studies should be repeated to confirm the diagnosis and rule out other pathogens before instituting another course of therapy.
Clindamycin is found in very low levels in the serum (5%) following multiple topical applications. No systemic accumulation is seen.
Intravaginal administration of clotrimazole in human has negligible systemic absorption (3-10%)
Thus with Tivag no systemic effects are anticipated.
Tivag is to be administered intravaginally and is not to be taken orally.

Pregnancy and Lactation: Experimental and clinical investigations given on indication of harmful effects to the mother or child when clotrimazole is used during pregnancy. However, there are no adequate and well controlled studies of use of clindamycin in pregnant women. Hence Tivag should be used during pregnancy only if it is a must (under the advice of physician).
Tivag may be used in lactation with caution.

Adverse Effect: Tivag vaginal capsules are well tolerated locally i.e. intravaginally. In isolated cases, vaginal irritation, itching, burning, dyspareunia and in rare cases slight increase in urinary frequency and burning or irritation intolerance and in such cases the treatment has to be stopped. Hypersensitivity reactions may occur.

Important: Only vaginal insertion. Not for oral use.
Pregnant, Lactating women should strictly abide by the instructions of their physicians.
Presentation (Blister Pack): Pack of 7 soft gelatin vaginal capsules.
Application and Dosage:
One capsule daily should be inserted into the vagina, for 7 consecutive nights preferably before retiring to bed. Treatment should be timed so as to avoids the menstrual period.
Directions for use:
1. *Wash hands thoroughly before and after insertion of Tivag in the vagina*
2. *The capsule should be inserted as deep into the vagina as possible. This is best achieved when lying on the back with legs pulled in a little towards the body.*
3. After insertion of Tivag, no activity should be done such as standing, walking running etc. hence Tivag should be inserted while retiring to bed.
4. For best results take the complete therapy for 7 days. Do not discontinue in the middle.
5. Discontinue only if severe irritation is experienced only after consulting the doctor.
6. If a particular dose is missed, administer immediately when recalled.

STORAGE CONDITION: Store below 25°C in a cool & dry place. Protect from direct sunlight, heat & moisture.

Mfg Lic. No: 82/UA/2017
Manufactured. by :
Renowed Life Sciences
WHO-GMP Certified Co.
Plot No. 10 & 11, Sector 6-B, IIE, SIDCUL,
Haridwar 249403 , Uttarakhand (INDIA)

https://trungtamthuoc.com/