

Status: Currently Official on 17-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-9D13739B-E267-4F9E-A49F-85C5E20E6838_1_en-US
DOI: https://doi.org/10.31003/USPNF_M84150_01_01
DOI Ref: Iu5hb

© 2025 USPC
Do not distribute

Tolmetin Sodium Capsules

DEFINITION

Tolmetin Sodium Capsules contain an amount of tolmetin sodium equivalent to NLT 93.0% and NMT 107.0% of the labeled amount of tolmetin ($C_{15}H_{15}NO_3$).

IDENTIFICATION

- **A. ULTRAVIOLET ABSORPTION**

Standard stock solution: 0.1 mg/mL of [USP Tolmetin Sodium RS](#) in methanol

Standard solution: 0.01 mg/mL of [USP Tolmetin Sodium RS](#) from the *Standard stock solution* diluted with 0.1 N sodium hydroxide

Sample solution: Transfer a quantity of Capsule contents, equivalent to 10 mg of tolmetin, to a 100-mL volumetric flask. Add 50 mL of methanol, shake for 2 min, and dilute with methanol to volume. Filter a portion of this solution, transfer 10.0 mL of the filtrate to a second 100-mL volumetric flask, and dilute with 0.1 N sodium hydroxide to volume.

Analysis

Samples: *Standard solution* and *Sample solution*

Concomitantly measure the UV absorption.

Acceptance criteria: The UV absorption spectrum of the *Sample solution* exhibits maxima and minima at the same wavelengths as that of the *Standard solution*.

• **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

- **PROCEDURE**

Buffer: 1.7 g/L of tetrabutylammonium phosphate. Adjust with phosphoric acid to a pH of 2.7 ± 0.1 .

Mobile phase: Acetonitrile and *Buffer* (36:64)

Diluent: Acetonitrile and 0.01 N sodium hydroxide (40:60)

System suitability solution: 200 μ g/mL of *p*-toluic acid and 500 μ g/mL of [USP Tolmetin Sodium RS](#) in *Diluent*

Standard solution: 0.65 mg/mL of anhydrous tolmetin sodium from [USP Tolmetin Sodium RS](#) in *Diluent*

Sample solution: Transfer the equivalent of 60 mg of tolmetin from the powdered contents of NLT 20 Capsules to a 100-mL volumetric flask. Add 75 mL of *Diluent*, and shake by mechanical means for 30 min. Dilute with *Diluent* to volume, and pass through a filter of 0.45- μ m or finer pore size.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 254 nm

Column: 4.6-mm \times 15-cm; 5- μ m packing L7

Column temperature: $40 \pm 1.0^\circ$

Flow rate: 3 mL/min

Injection volume: 20 μ L

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 1.2 between the *p*-toluic acid and tolmetin sodium peaks, *System suitability solution*

Relative standard deviation: NMT 3.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of tolmetin ($C_{15}H_{15}NO_3$) in the portion of Capsule contents taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of the [USP Tolmetin Sodium RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of tolmetin in the *Sample solution* (mg/mL)

M_{r1} = molecular weight of tolmetin, 257.29

M_{r2} = molecular weight of anhydrous tolmetin sodium, 279.27

Acceptance criteria: 93.0%–107.0%

PERFORMANCE TESTS

- [DISSOLUTION \(711\)](#).

Solution A: Dissolve 2.0 g of sodium chloride in 7.0 mL of hydrochloric acid, and add water to 1000 mL.

Solution B: Dissolve 6.8 g of monobasic potassium phosphate in 250 mL of water, and add 190 mL of 0.2 N sodium hydroxide and 400 mL of water. Adjust with 0.2 N sodium hydroxide to a pH of 7.4–7.6. Dilute with water to 1000 mL.

Medium: *Solution A* and *Solution B* (336:664). Adjust the solution with small amounts of either solution to a pH of 4.5; 900 mL.

Apparatus 2: 50 rpm

Time: 30 min

Standard solution: [USP Tolmetin Sodium RS](#) in *Medium* at a known concentration

Sample solutions: Sample per [Dissolution \(711\)](#). Dilute with 0.1 N sodium hydroxide, as needed, and filter.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 322 nm

Analysis

Samples: *Standard solution* and *Sample solution*

Determine the quantity of tolmetin ($C_{15}H_{15}NO_3$) dissolved using filtered portions of the solution under test, suitably diluted with 0.1 N sodium hydroxide, in comparison with the *Standard solution*.

Tolerances: NLT 85% (Q) of the labeled amount of tolmetin ($C_{15}H_{15}NO_3$) is dissolved.

- [UNIFORMITY OF DOSAGE UNITS \(905\)](#): Meet the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers.

- [USP REFERENCE STANDARDS \(11\)](#).

[USP Tolmetin Sodium RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
TOLMETIN SODIUM CAPSULES	Documentary Standards Support	SM22020 Small Molecules 2
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM22020 Small Molecules 2

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. 46(5)

Current DocID: [GUID-9D13739B-E267-4F9E-A49F-85C5E20E6838_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_M84150_01_01

OFFICIAL