

Status: Currently Official on 17-Feb-2025  
 Official Date: Official as of 01-May-2024  
 Document Type: USP Monographs  
 DocId: GUID-E42D32D5-DD86-47D8-A822-59CAF3F4792\_2\_en-US  
 DOI: [https://doi.org/10.31003/USPNF\\_M84240\\_02\\_01](https://doi.org/10.31003/USPNF_M84240_02_01)  
 DOI Ref: g65tw

© 2025 USPC  
 Do not distribute

# Tolnaftate Topical Powder

## DEFINITION

Tolnaftate Topical Powder contains NLT 90.0% and NMT 110.0% of the labeled amount of tolinaftate (C<sub>19</sub>H<sub>17</sub>NOS).

## IDENTIFICATION

### Change to read:

- ▲ **A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.▲ (USP 1-May-2024)

### Add the following:

- ▲ **B.** The UV spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.▲ (USP 1-May-2024)

## ASSAY

### Change to read:

#### PROCEDURE

▲ **Solution A:** [Methanol](#) and [water](#) (70:30). To each liter of the solution, add 1 mL of [trifluoroacetic acid](#).

**Solution B:** [Methanol](#) and [water](#) (90:10). To each liter of the solution, add 1 mL of [trifluoroacetic acid](#).

**Mobile phase:** See [Table 1](#).

Table 1

Time (min)	Solution A (%)	Solution B (%)
0	100	0
12	100	0
30	0	100
33	0	100

**Standard solution:** 0.05 mg/mL of [USP Tolnaftate RS](#) in [methanol](#)

**Sample solution:** Nominally 0.05 mg/mL of tolinaftate prepared as follows. Transfer a quantity of the Topical Powder equivalent of 2.5 mg of tolinaftate into a 50-mL volumetric flask, add 30 mL of [methanol](#), shake for 1 h, and dilute with [methanol](#) to volume. Centrifuge at 8000 rpm for 10 min, pass a portion of the supernatant through a suitable filter of 0.45-µm pore size, and discard the first 2 mL.

## Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** LC

**Detector:** UV 254 nm. For *Identification B*, use a diode array detector in the range of 200–400 nm.

**Column:** 4.6-mm × 15-cm; 5-µm packing [L1](#)

### Temperatures

**Autosampler:** 4°

**Column:** 30°

**Flow rate:** 1 mL/min

**Injection volume:** 10 µL

**System suitability**

**Sample:** *Standard solution*

**Suitability requirements**

**Tailing factor:** NMT 2.0

**Relative standard deviation:** NMT 1.0%

**Analysis**

**Samples:** *Standard solution and Sample solution*

Calculate the percentage of the labeled amount of tolnaftate (C<sub>19</sub>H<sub>17</sub>NOS) in the portion of the Topical Powder taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

*r<sub>U</sub>* = peak response of tolnaftate from the *Sample solution*

*r<sub>S</sub>* = peak response of tolnaftate from the *Standard solution*

*C<sub>S</sub>* = concentration of [USP Tolnaftate RS](#) in the *Standard solution* (mg/mL)

*C<sub>U</sub>* = nominal concentration of tolnaftate in the *Sample solution* (mg/mL)▲ (USP 1-May-2024)

**Acceptance criteria:** 90.0%–110.0%

**PERFORMANCE TESTS**

- [MINIMUM FILL \(755\)](#): Meets the requirements

**ADDITIONAL REQUIREMENTS**

**Change to read:**

- **PACKAGING AND STORAGE:** Preserve in tight containers.▲Store between 2° and 30°.▲ (USP 1-May-2024)
- [USP REFERENCE STANDARDS \(11\)](#).  
[USP Tolnaftate RS](#)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
TOLNAFTATE TOPICAL POWDER	<a href="#">Documentary Standards Support</a>	SM12020 Small Molecules 1
REFERENCE STANDARD SUPPORT	RS Technical Services <a href="mailto:RSTECH@usp.org">RSTECH@usp.org</a>	SM12020 Small Molecules 1

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. 48(4)

**Current DocID:** GUID-E42D32D5-DD86-47D8-A822-59CAFC3F4792\_2\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_M84240\\_02\\_01](https://doi.org/10.31003/USPNF_M84240_02_01)

**DOI ref:** [g65tw](#)